

Clinical Pharmacogenetics Implementation Consortium (CPIC) Guideline for *CYP2C19* and Proton Pump Inhibitor Dosing

John J. Lima¹, Cameron D. Thomas², Julia Barbarino³, Zeruesenay Desta⁴, Sara L. Van Driest⁵, Nihal El Rouby^{2,6}, Julie A. Johnson², Larisa H. Cavallari², Valentina Shakhnovich^{7,8,9}, David L. Thacker^{10,11}, Stuart A. Scott^{12,13}, Matthias Schwab^{14,15,16}, Chakradhara Rao S Uppugunduri^{17,18}, Christine M. Formea¹⁹, James P. Franciosi^{20,21}, Katrin Sangkuhl³, Andrea Gaedigk⁷, Teri E. Klein³, Roseann S. Gammal^{22,23}, Takahisa Furuta²⁴

¹ Center for Pharmacogenomics and Translational Research, Nemours Children's Health, Jacksonville, FL, USA

² Department of Pharmacotherapy and Translational Research, and Center for Pharmacogenomics and Precision Medicine, College of Pharmacy, University of Florida, Gainesville, FL, USA

³ Department of Biomedical Data Science, Stanford University, Stanford, CA, USA

⁴ Department of Medicine, Division of Clinical Pharmacology, Indiana University School of Medicine, Indianapolis, IN, USA

⁵ Departments of Pediatrics and Medicine, Vanderbilt University School of Medicine, Nashville, TN, USA

⁶ Division of Pharmacy Practice & Administrative Sciences, University of Cincinnati James Winkle College of Pharmacy, Cincinnati, OH, USA

⁷ Division of Clinical Pharmacology, Toxicology & Therapeutic Innovation, Children's Mercy Kansas City and University of Missouri Kansas City School of Medicine, Kansas City, MO, USA

⁸ Division of Gastroenterology, Hepatology, and Nutrition, Children's Mercy Kansas City, Kansas City, MO, USA

⁹ Center for Children's Healthy Lifestyles & Nutrition, Kansas City, MO, USA

¹⁰ Department of Clinical Pharmacology, Indiana University School of Medicine, Indianapolis, IN, USA

¹¹ Translational Software, Bellevue, WA, USA

¹² Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA

¹³ Sema4, Stamford, CT, USA

¹⁴ Dr Margarete Fischer-Bosch-Institute of Clinical Pharmacology, Stuttgart, Germany

¹⁵ Department of Clinical Pharmacology, University Hospital, Tuebingen, Germany

¹⁶ Department of Pharmacy and Biochemistry, University of Tuebingen, Tuebingen, Germany

¹⁷ CANSEARCH Research Laboratory, Department of Paediatrics, Gynaecology and Obstetrics, Faculty of Medicine, University of Geneva, Geneva, Switzerland

¹⁸ Oncology-Hematology Unit, Department of Paediatrics, Gynaecology and Obstetrics, Geneva University Hospital, Geneva, Switzerland

¹⁹ Department of Pharmacy Services and Intermountain Precision Genomics, Intermountain Healthcare, Salt Lake City, UT, USA

²⁰ Division of Gastroenterology, Hepatology and Nutrition, Nemours Children's Hospital, Orlando, FL

²¹ Department of Pediatrics, University of Central Florida College of Medicine, Orlando, FL, USA

²² Department of Pharmacy Practice, MCPHS University School of Pharmacy, Boston, MA, USA

²³ Department of Pharmaceutical Sciences, St. Jude Children's Research Hospital, Memphis, TN, USA

²⁴ Center for Clinical Research, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka, Japan

Corresponding Author:

John J. Lima, PharmD
Principal Research Scientist Emeritus
Center for Pharmacogenomics and Translational Research
Nemours Children's Health
807 Children's Way
Jacksonville, FL 32207
Phone: 352-219-3048
Fax: 904-697-3799
Email: john.lima@nemours.org; contact@cpicpgx.org

Word Counts:

Abstract (250 limit): 153
Text (4,000 limit): 3,435
References (50 limit): 39
Figures/tables (7 limit): 2

Keywords: CYP2C19, PPI, proton pump inhibitor, omeprazole, lansoprazole, pantoprazole, dexlansoprazole, esomeprazole, rabeprazole, pharmacogenetics, pharmacogenomics, CPIC, metabolism

Conflicts of Interest:

JAJ is a consultant to United Health Group for their plans to begin implementing pharmacogenomics in clinical practice; SAS is a paid employee of Sema4, which is a for-profit genetic testing company that offers pharmacogenetic testing. DLT is a paid employee of Translational Software, which is a for-profit genetic testing company that offers pharmacogenetic testing. As an Associate Editor for *Clinical Pharmacology & Therapeutics*, SLV was not involved in the review or decision process for this paper. All other authors declared no competing interests for this work.

Funding:

This work was funded by the National Institutes of Health (NIH) for CPIC (R24GM115264 and U24HG010135), PharmGKB (U24HG010615) and PharmVar (R24 GM123930). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. Additional grant funding includes NIH IGNITE Network (NIH U01 HG007269) (JAJ, JJJ, LHC) and U01 HG007762, R01 GM121707 and R01 GM078501 (ZD); GM 0NIH-NCATS UL1TR001427 (LHC); CANSEARCH foundation, Switzerland (CRU); Robert Bosch Stiftung, Stuttgart, Germany (MS); Innovation in Regulatory Science Award 1015006 from the Burroughs Wellcome Fund (SLV); 1K23DK115827-01A1 and the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition Foundation (VS); and the EU H2020 UPGx grant 668353 (MS).

ABSTRACT

Proton pump inhibitors (PPIs) are widely used for acid suppression in the treatment and prevention of many conditions including gastroesophageal reflux disease, gastric and duodenal ulcers, erosive esophagitis, *H. pylori* infection, and pathological hypersecretory conditions. Most PPIs are metabolized primarily by CYP2C19 into inactive metabolites, and *CYP2C19* genotype has been linked to PPI exposure, efficacy, and adverse effects. We summarize the evidence from the literature and provide therapeutic recommendations for PPI prescribing based on *CYP2C19* genotype (updates at www.cpicpgx.org). The potential benefits of using *CYP2C19* genotype data to guide PPI therapy include 1) identifying patients with genotypes predictive of lower plasma exposure and prescribing them a higher dose that will increase the likelihood of efficacy; and 2) identifying patients on chronic therapy with genotypes predictive of higher plasma exposure and prescribing them a decreased dose to minimize the risk of toxicity that is associated with long-term PPI use, particularly at higher plasma concentrations.

INTRODUCTION

Proton pump inhibitors (PPIs) are widely used for acid suppression in the treatment and prevention of a variety of conditions including gastroesophageal reflux disease (GERD), gastric and duodenal ulcers, erosive esophagitis, eosinophilic esophagitis, *Helicobacter pylori* (*H. pylori*) infection, and pathological hypersecretory conditions in adults and children. Most PPIs are extensively metabolized into inactive metabolites primarily by the hepatic cytochrome P450 2C19 (CYP2C19) enzyme, and *CYP2C19* genotypes have been linked to PPI exposure, with lower exposure associated with treatment failure and higher exposure associated with improved efficacy (1). Higher exposure of PPIs has also been associated with adverse effects (1), as has long-term use (2). The purpose of this guideline is to provide clinicians with information that facilitates the interpretation of clinical *CYP2C19* genotyping test results to guide PPI prescribing. Detailed guidelines for the use of PPIs, cost effectiveness of *CYP2C19* genotyping, and whether to order a *CYP2C19* genotype test prior to PPI prescribing, are beyond the scope of this document. Clinical Pharmacogenetics Implementation Consortium (CPIC®) guidelines are periodically updated at www.cpicpgx.org/guidelines/.

FOCUSED LITERATURE REVIEW

A systematic literature review focused on the link between *CYP2C19* genotypes and PPI metabolism, exposure, efficacy and adverse effects was conducted (details in **Supplemental Material**).

GENE: *CYP2C19*

Background

The *CYP2C19* gene is highly polymorphic with 37 known variant star (*) alleles, including rare copy number variants (i.e., gene deletions) (<https://www.pharmvar.org/gene/CYP2C19>; see ***CYP2C19* Allele Definition Table** online (3, 4)). The frequencies of these alleles significantly differ across ancestrally diverse populations (see ***CYP2C19* Frequency Table** online (3, 4)). Alleles are categorized into functional groups as follows: normal function (e.g., *CYP2C19**1), decreased function (e.g., *CYP2C19**9), no function (e.g., *CYP2C19**2 and *3), and increased function (e.g., *CYP2C19**17). Clinical allele function, as described in the ***CYP2C19* Allele Functionality Table**, was determined based on reported *in vitro* and/or *in vivo* data when available (3, 4). The most common *CYP2C19* no function allele, *CYP2C19**2 (c.681G>A; rs4244285), has an allele frequency of ~15% in Europeans and Africans, ~25-30% in Asians, and ~60% in Oceanians. Other *CYP2C19* variant alleles with decreased or no function (e.g., *3-*8) typically have allele frequencies below 1%, with the exception of *CYP2C19**3 (c.636G>A; rs4986893) in Asians (allele frequency of 2-7%) and Oceanians (allele frequency of 15%). The increased function allele *CYP2C19**17 (c.-806C>T; rs12248560) is most common in African, European, and Near Eastern populations, with an allele frequency of ~20%.

Genetic Test Interpretation

The combination of inherited alleles determines a person's diplotype (also referred to as genotype). **Table 1** defines each predicted phenotype based on allele function combinations and provides example diplotypes. *CYP2C19* normal metabolizers (NMs) are characterized by the presence of two normal function alleles (e.g., *CYP2C19* *1/*1). *CYP2C19* intermediate metabolizers (IMs) are characterized by the presence of one normal function allele and one no

function allele (e.g., *CYP2C19* *1/*2), or one no function allele and one increased function allele (e.g., *CYP2C19* *2/*17). Limited data suggest that the increased function allele *CYP2C19**17 may not compensate for no function alleles such as *CYP2C19**2 (5). *CYP2C19* poor metabolizers (PMs) are characterized by the presence of two no function alleles (e.g., *CYP2C19* *2/*2). Diplotypes characterized by one normal function allele and one increased function allele (i.e., *CYP2C19* *1/*17) are classified as rapid metabolizers (RMs), and diplotypes characterized by two increased function alleles (i.e., *CYP2C19* *17/*17) are classified as ultrarapid metabolizers (UMs). There are limited data available for decreased function alleles (e.g., *CYP2C19**9); therefore, individuals who have one normal function and one decreased function allele, or one increased function and one decreased function allele, or two decreased function alleles, are currently classified as “likely IM.” Individuals with one no function and one decreased function allele are currently classified as “likely PM.” The “indeterminate” phenotype is assigned when the individual carries one or two uncertain function alleles. See the ***CYP2C19* Diplotype-Phenotype Table** online for a complete list of possible diplotypes and the corresponding predicted phenotype assignments (3, 4).

Clinical laboratories report *CYP2C19* genotype results using star (*) allele nomenclature. The star (*) allele nomenclature for *CYP2C19* is found at the Pharmacogene Variation (PharmVar) Consortium website (<https://www.pharmvar.org/gene/CYP2C19>). Tables on the CPIC website contain a list of *CYP2C19* alleles, the combinations of variants that define each allele, allele clinical functional status, and reported allele frequencies across major ancestral populations (3).

Available Genetic Test Options

See the Genetic Testing Registry (www.ncbi.nlm.nih.gov/gtr/) for more information on commercially available clinical testing options.

Incidental Findings

No inherited diseases or conditions have been consistently or strongly linked to germline genetic variants in *CYP2C19* independent of drug metabolism and response.

Other Considerations

Not applicable

DRUGS: PROTON PUMP INHIBITORS

Background

PPIs are substituted benzimidazoles that inhibit the final pathway of acid production in gastric parietal cells by covalently binding to the H^+/K^+ ATPase proton pump, which leads to inhibition of gastric acid secretion that lasts for 24-48 hours despite short PPI half-lives (1, 6). This irreversible inhibition is only overcome by synthesis of new H^+/K^+ ATPase proton pumps, which may take ~54 hours to fully regenerate (6). Six PPIs are currently used in clinical practice including the first-generation inhibitors omeprazole, lansoprazole and pantoprazole, and the second-generation inhibitors esomeprazole, rabeprazole, and dexlansoprazole. CYP2C19 is a major metabolic pathway for the clearance of first generation PPIs (~80%) with a lesser contribution by CYP3A4. Dexlansoprazole (R-lansoprazole) appears to share a similar metabolic pathway to lansoprazole. In contrast, the second generation PPIs esomeprazole and rabeprazole are less dependent on CYP2C19 in their metabolism, suggesting that they may be less influenced

by genetic variability in *CYP2C19* compared to first generation PPIs. Specifically, rabeprazole is primarily cleared by non-enzymatic mechanisms (1). Omeprazole and esomeprazole exhibit nonlinear pharmacokinetics due to *CYP2C19* autoinhibition leading to an increased area under the serum concentration versus time curve (AUC) with repeated administration. The elevation in AUC that results from autoinhibition is greater for esomeprazole (1.45-1.74 fold) than omeprazole (7).

While PPIs have been among the most commonly prescribed medications due in part to the perception that they have a high safety to risk profile, a large body of evidence is emerging that links adverse events with long-term PPI use (2). PPI use has been associated with numerous adverse events, including electrolyte imbalances (e.g., hypomagnesemia), infections, kidney disease, and bone fractures (1).

Linking Genetic Variability to Variability in Drug-related Phenotypes

First Generation PPIs: Omeprazole, Lansoprazole, and Pantoprazole

There is a substantial body of evidence linking *CYP2C19* genotype with variability in plasma concentrations and efficacy of first-generation PPIs (omeprazole, lansoprazole, and pantoprazole). As outlined in **Tables S1-S3**, the evidence associating *CYP2C19* genotype with omeprazole, lansoprazole, and pantoprazole plasma concentrations was graded as high. Multiple studies have shown that the *CYP2C19* IM and PM phenotypes are associated with decreased clearance and increased plasma concentrations of these PPIs leading to increased treatment success compared to *CYP2C19* NMs, including for *H. pylori* infection and erosive esophagitis (8-10). In contrast, *CYP2C19* RMs and UMs have increased PPI clearance and decreased plasma concentrations compared to *CYP2C19* NMs, which may increase risk of treatment failure

compared to CYP2C19 NMs, IMs, and PMs (11, 12). It is important to note that most *CYP2C19* studies evaluating PPIs were conducted in Asian populations, in whom the frequency of the increased function *CYP2C19*17* allele is low compared to non-Asians; therefore, few studies including CYP2C19 RMs and UMs have been published to date. Prescribing recommendations for CYP2C19 RMs and UMs in this guideline were based on pharmacokinetic differences versus NMs and differences in PPI effectiveness between NMs and IMs/PMs. This body of literature provides the basis for the prescribing recommendations presented in **Table 2**.

Second Generation PPIs: Esomeprazole, Rabeprazole, and Dexlansoprazole

There is less evidence linking *CYP2C19* genotype with variability in plasma concentrations and effectiveness of second-generation PPIs (esomeprazole, rabeprazole, and dexlansoprazole), both in terms of number of studies and strength of the association. As outlined in **Table S4-S6**, the evidence associating *CYP2C19* genotype with esomeprazole, rabeprazole, and dexlansoprazole plasma concentrations, efficacy, and toxicity was graded as moderate or weak. While less data exist on the influence of *CYP2C19* genotype on dexlansoprazole compared to first generation PPIs, similar effects of *CYP2C19* genotype on dexlansoprazole pharmacokinetics and effectiveness are expected given its similar metabolic pathway as lansoprazole (1, 13). Inconsistent findings regarding the effect of *CYP2C19* genotype on the pharmacokinetics and therapeutic response to esomeprazole and rabeprazole preclude making recommendations for these second-generation PPIs (i.e., CPIC level C; no recommendation).

Therapeutic Recommendations

Table 2 summarizes therapeutic recommendations for PPI prescribing in adults and pediatric patients based on CYP2C19 phenotype, specifically for the first-generation PPIs (omeprazole, lansoprazole, pantoprazole) and dexlansoprazole. These recommendations apply to both oral and intravenous PPI use. While CYP2C19 NMs are expected to have normal PPI metabolism and clearance, a large body of literature from studies in Asian populations reported an association between CYP2C19 NMs and decreased therapeutic effectiveness with these PPIs (e.g., failure to eradicate *H. pylori* infection and lower healing rates of erosive esophagitis) compared to CYP2C19 IMs and PMs (**Tables S1-S4**). Therefore, for CYP2C19 NMs, initiating these PPIs at standard daily doses (e.g., label recommended doses) is generally recommended; however, for *H. pylori* infection or erosive esophagitis, clinicians may consider increasing the recommended dose for these indications by 50-100% to optimize therapeutic efficacy.

Following administration of standard doses of first generation PPIs, CYP2C19 IMs and PMs experience higher PPI AUC (3-14 fold) and Cmax (2-6 fold) compared to CYP2C19 NMs as a result of reduced PPI clearance via the CYP2C19 pathway (14-18). The increased PPI exposure in CYP2C19 IMs and PMs has been linked to improved acid suppression (i.e., higher intragastric pH and longer time with pH > 4.0) and improved therapeutic benefits. Thus, CYP2C19 IMs and PMs are considered to be “therapeutically advantaged” compared to NMs in terms of efficacy (19-23). However, it has been suggested that continued inhibition of acid secretion in individuals taking PPIs chronically who are genotyped as CYP2C19 IMs or PMs may have a higher risk of PPI-related adverse events compared to NM, RM, or UM phenotypes (1). While the current data are insufficient to make strong dosing recommendations, potential associations of CYP2C19 phenotype and incidence of adverse events (e.g., infections) are emerging (24). Therefore, for CYP2C19 IMs and PMs, it is recommended to initiate standard

daily dosing to maximize the likelihood of efficacy and, once efficacy is achieved, consider a 50% reduction in the daily dose in the setting of chronic PPI therapy (beyond 12 weeks) to minimize the risk of adverse events from prolonged acid suppression. If a dose reduction is made, monitoring for continued efficacy is recommended. Additional studies that investigate the relationship between *CYP2C19* genotype and incidence of PPI-related adverse events are needed.

The RM and UM phenotypes are driven by the presence of the increased function *CYP2C19*17* allele. Due to the relatively recent discovery of this variant (11) and because the majority of studies describing associations between *CYP2C19* genotype, pharmacokinetics, and pharmacodynamics of PPIs were conducted in Asian populations in whom the *CYP2C19*17* allele occurs less frequently, there are limited data on the relationship between *CYP2C19*17*, pharmacokinetic parameters, acid secretion indices and therapeutic outcomes in *CYP2C19* RMs and UMs. Additional studies with *CYP2C19* RMs and UMs are needed. Nevertheless, the low PPI exposure documented in patients who are *CYP2C19* UMs compared to NMs, IMs, and PMs suggests that these individuals may benefit from higher than standard daily doses of PPIs (**Tables S1-S3**). Therefore, it is recommended to increase the starting daily dose by 100% in *CYP2C19* UMs. For RMs, standard dosing should be initiated, but a 50-100% dose increase could be considered for the treatment of *H. pylori* infection and erosive esophagitis to maximize the likelihood of therapeutic plasma concentrations and therapeutic effect. These patients should be monitored for efficacy.

The plasma half-life of PPIs is short (~30 minutes to 5 hours), but the biological effects they exert is much longer, as it takes ~54 hours to regenerate new acid pumps after inactivation by PPIs. Studies have documented that daily doses administered two to four times daily may result in improved efficacy compared to the same total daily dose given once daily (25, 26).

Although adherence to PPI dosing three to four times per day to overcome the short half-life may be challenging, it is recommended that increased PPI doses (50-100%) be administered as twice daily dosing, and more frequent dosing intervals could be considered for increased benefit, with the caveat that this dosing regimen may compromise compliance.

There are fewer data available investigating the association between dexlansoprazole and CYP2C19 metabolizer status compared to the first-generation PPIs. However, reported pharmacokinetic data support the association between CYP2C19 IMs and PMs and increased dexlansoprazole exposure (13, 27). Additionally, CYP2C19 PMs were reported to have greater acid suppression compared to NMs (13). Given the similarity in metabolism between lansoprazole and dexlansoprazole, it is reasonable to extrapolate the recommendations from the first-generation PPIs (**Table 2**). These recommendations are considered “optional” due to the limited data with dexlansoprazole and *CYP2C19*.

Pediatrics

The *CYP2C19*-guided PPI recommendations presented in **Table 2** also apply to pediatric patients. PPI use in children is common and continues to increase. PPIs have U.S. Food and Drug Administration-approved indications in children for the short-term treatment of symptomatic GERD, healing of erosive esophagitis, treatment of peptic ulcer disease, and eradication of *H. pylori*. PPIs are also considered standard of care for pediatric eosinophilic esophagitis. Off-label and potentially inappropriate use of long-term PPI therapy in children is common, particularly in infants less than one year of age for uncomplicated, physiologic, gastroesophageal reflux and colic (28).

In children older than one year of age, there is emerging evidence that *CYP2C19* genetic variation influences PPI pharmacokinetics and response (24, 29-31). *CYP2C19* RM and UM phenotypes have been associated with decreased efficacy compared to PM and NM phenotypes when treating pediatric GERD and eosinophilic esophagitis (32-34). The *CYP2C19* PM phenotype is associated with higher rates of respiratory and gastrointestinal infections than the NM, RM, or UM phenotypes (35). A recent pilot study of *CYP2C19*-genotype-guided dosing of PPIs in children has been promising, and additional studies are ongoing (36, 37). These reports support genotype-based optimization of PPI therapy for children. However, very low clearance in pre-term infants and infants less than 2-3 months of age (29) makes recommendations in the neonatal population difficult to support. Additional pediatric considerations are discussed in the **Supplemental Material**.

Recommendations for Incidental Findings

Not applicable

Other Considerations

Drug-Drug-Gene Interactions

In addition to *CYP2C19*, most PPIs are metabolized to some extent by *CYP3A*, and the fraction of metabolism by this alternative pathway increases in *CYP2C19* IMs and PMs (38). Due to drug-drug-gene interactions, concomitant administration of strong inhibitors of *CYP3A* may increase the risk for adverse effects during chronic dosing with PPIs in *CYP2C19* IMs and PMs. In addition, strong inhibitors of *CYP2C19* (e.g., fluvoxamine) can lead to phenoconversion that substantially increases systemic exposure to PPIs (except in *CYP2C19* PMs), while inducers

of CYP2C19 and CYP3A (e.g., rifampin) can lead to reduced exposure and treatment failure. Therefore, assessment of drug interactions may be needed when a PPI and a perpetrator drug are co-administered chronically.

Implementation of this Guideline

The guideline supplement and CPIC website (<https://cpicpgx.org/guidelines/cpic-guideline-for-proton-pump-inhibitors-and-cyp2c19/>) contains resources that can be used within electronic health records (EHRs) to assist clinicians in applying genetic information to patient care for the purpose of drug therapy optimization (see *Resources to incorporate pharmacogenetics into an electronic health record with clinical decision support* in the **Supplemental Material**).

POTENTIAL BENEFITS AND RISKS FOR THE PATIENT

The potential benefit of using *CYP2C19* genotype data to guide PPI therapy is that patients with genotypes predictive of lower plasma exposure may be identified and prescribed an increased dose that will increase the likelihood of efficacy. Patients on chronic therapy with genotypes predictive of higher plasma exposure may consider a dose reduction to minimize the risk of toxicity that is associated with long-term PPI use (over exposure), particularly at higher plasma concentrations. The potential risks of genotype-guided PPI therapy include therapeutic failure in patients for whom a dose decrease was recommended and increased risk of toxicity in patients for whom a dose increase was recommended. As with any laboratory test, a possible risk to patients is an error in genotyping or phenotype prediction, which could have long-term adverse health implications for patients.

CAVEATS: APPROPRIATE USE AND/OR POTENTIAL MISUSE OF GENETIC TESTS

There are some important limitations to *CYP2C19* genetic tests. Targeted genotyping tests focus on interrogating previously described star (*) alleles and therefore are not designed to detect novel variants. Furthermore, rare allelic *CYP2C19* variants may not be included in the genotype test used, and patients with these rare variants may be assigned a NM phenotype (*CYP2C19**1/*1) by default. As such, an assigned *1 allele could potentially harbor an undetected *CYP2C19* genetic variant that results in altered metabolism and drug exposure. In addition, rare alleles with gene deletions at the *CYP2C19* locus have recently been reported (*36 and *37) (39); however, most clinical laboratories do not currently test for *CYP2C19* copy number variants or deletions. Therefore, it is important that clinical providers appreciate the limitations of targeted genotyping tests and understand which *CYP2C19* variant alleles were genotyped when interpreting results. As with any diagnostic test, *CYP2C19* genotype is just one factor that clinicians should consider when prescribing PPIs.

DISCLAIMER

Clinical Pharmacogenetics Implementation Consortium (CPIC) guidelines reflect expert consensus based on clinical evidence and peer-reviewed literature available at the time they are written and are intended only to assist clinicians in decision-making, as well as to identify questions for further research. New evidence may have emerged since the time a guideline was submitted for publication. Guidelines are limited in scope and are not applicable to interventions or diseases not specifically identified. Guidelines do not account for all individual variation

among patients and cannot be considered inclusive of all proper methods of care or exclusive of other treatments. It remains the responsibility of the health care provider to determine the best course of treatment for the patient. Adherence to any guideline is voluntary, with the ultimate determination regarding its application to be solely made by the clinician and the patient. CPIC assumes no responsibility for any injury to persons or damage to property related to any use of CPIC's guidelines, or for any errors or omissions.

ACKNOWLEDGEMENTS

We acknowledge the critical input of Dr. Mary Relling, Dr. Kelly Caudle, and members of the Clinical Pharmacogenetics Implementation Consortium (CPIC) of the Pharmacogenomics Research Network, funded by the National Institutes of Health. CPIC members are listed here:

<https://cpicpgx.org/members/>.

REFERENCES

- (1) El Rouby, N., Lima, J.J. & Johnson, J.A. Proton pump inhibitors: from CYP2C19 pharmacogenetics to precision medicine. *Expert Opin Drug Metab Toxicol* **14**, 447-60 (2018).
- (2) Jaynes, M. & Kumar, A.B. The risks of long-term use of proton pump inhibitors: a critical review. *Ther Adv Drug Saf* **10**, 2042098618809927 (2019). doi: 10.1177/2042098618809927.
- (3) CPIC. *CPIC® Guideline for Proton Pump Inhibitors and CYP2C19*. <<https://cpicpgx.org/guidelines/cpic-guideline-for-proton-pump-inhibitors-and-cyp2c19/>> (2020).
- (4) PharmGKB. *Gene-specific Information Tables for CYP2C19*. <<https://www.pharmgkb.org/page/cyp2c19RefMaterials>>. Accessed Jan 20 2020.
- (5) Sibbing, D. *et al.* Isolated and interactive impact of common CYP2C19 genetic variants on the antiplatelet effect of chronic clopidogrel therapy. *J Thromb Haemost* **8**, 1685-93 (2010).
- (6) Shin, J.M. & Sachs, G. Pharmacology of proton pump inhibitors. *Curr Gastroenterol Rep* **10**, 528-34 (2008).
- (7) Hassan-Alin, M., Andersson, T., Niazi, M. & Rohss, K. A pharmacokinetic study comparing single and repeated oral doses of 20 mg and 40 mg omeprazole and its two

- optical isomers, S-omeprazole (esomeprazole) and R-omeprazole, in healthy subjects. *Eur J Clin Pharmacol* **60**, 779-84 (2005).
- (8) Furuta, T. *et al.* Effect of genetic differences in omeprazole metabolism on cure rates for *Helicobacter pylori* infection and peptic ulcer. *Ann Intern Med* **129**, 1027-30 (1998).
 - (9) Ichikawa, H., Sugimoto, M., Sugimoto, K., Andoh, A. & Furuta, T. Rapid metabolizer genotype of CYP2C19 is a risk factor of being refractory to proton pump inhibitor therapy for reflux esophagitis. *J Gastroenterol Hepatol* **31**, 716-26 (2016).
 - (10) Lin, Y.A. *et al.* Effect of CYP2C19 Gene Polymorphisms on Proton Pump Inhibitor, Amoxicillin, and Levofloxacin Triple Therapy for Eradication of *Helicobacter Pylori*. *Med Sci Monit* **23**, 2701-7 (2017).
 - (11) Sim, S.C. *et al.* A common novel CYP2C19 gene variant causes ultrarapid drug metabolism relevant for the drug response to proton pump inhibitors and antidepressants. *Clin Pharmacol Ther* **79**, 103-13 (2006).
 - (12) Gawronska-Szklarz, B. *et al.* CYP2C19 polymorphism affects single-dose pharmacokinetics of oral pantoprazole in healthy volunteers. *Eur J Clin Pharmacol* **68**, 1267-74 (2012).
 - (13) Sun, L.N. *et al.* Impact of Gastric H(+)/K(+)-ATPase rs2733743 on the Intragastric pH-Values of Dexamprazole Injection in Chinese Subjects. *Front Pharmacol* **8**, 670 (2017).
 - (14) Chang, M. *et al.* Interphenotype differences in disposition and effect on gastrin levels of omeprazole--suitability of omeprazole as a probe for CYP2C19. *Br J Clin Pharmacol* **39**, 511-8 (1995).
 - (15) Tanaka, M. *et al.* Stereoselective pharmacokinetics of pantoprazole, a proton pump inhibitor, in extensive and poor metabolizers of S-mephenytoin. *Clin Pharmacol Ther* **69**, 108-13 (2001).
 - (16) Kim, K.A. *et al.* Enantioselective disposition of lansoprazole in extensive and poor metabolizers of CYP2C19. *Clin Pharmacol Ther* **72**, 90-9 (2002).
 - (17) He, N. *et al.* Inhibitory effect of troleandomycin on the metabolism of omeprazole is CYP2C19 genotype-dependent. *Xenobiotica* **33**, 211-21 (2003).
 - (18) Qiao, H.L. *et al.* Pharmacokinetics of three proton pump inhibitors in Chinese subjects in relation to the CYP2C19 genotype. *Eur J Clin Pharmacol* **62**, 107-12 (2006).
 - (19) Furuta, T. *et al.* CYP2C19 genotype status and effect of omeprazole on intragastric pH in humans. *Clin Pharmacol Ther* **65**, 552-61 (1999).
 - (20) Shimatani, T., Inoue, M., Kuroiwa, T., Horikawa, Y., Mieno, H. & Nakamura, M. Effect of omeprazole 10 mg on intragastric pH in three different CYP2C19 genotypes, compared with omeprazole 20 mg and lafutidine 20 mg, a new H₂-receptor antagonist. *Aliment Pharmacol Ther* **18**, 1149-57 (2003).
 - (21) Park, S. *et al.* Effects of CYP2C19 Genetic Polymorphisms on PK/PD Responses of Omeprazole in Korean Healthy Volunteers. *J Korean Med Sci* **32**, 729-36 (2017).
 - (22) Chen, W.Y., Chang, W.L., Tsai, Y.C., Cheng, H.C., Lu, C.C. & Sheu, B.S. Double-dosed pantoprazole accelerates the sustained symptomatic response in overweight and obese patients with reflux esophagitis in Los Angeles grades A and B. *Am J Gastroenterol* **105**, 1046-52 (2010).
 - (23) Kurzawski, M., Gawronska-Szklarz, B., Wrzesniewska, J., Siuda, A., Starzynska, T. & Drozdziak, M. Effect of CYP2C19*17 gene variant on *Helicobacter pylori* eradication in peptic ulcer patients. *Eur J Clin Pharmacol* **62**, 877-80 (2006).

- (24) Bernal, C.J. *et al.* CYP2C19 Phenotype and Risk of Proton Pump Inhibitor-Associated Infections. *Pediatrics* **144**, (2019). doi: 10.1542/peds.2019-0857.
- (25) Furuta, T. *et al.* Pharmacogenomics-based tailored versus standard therapeutic regimen for eradication of *H. pylori*. *Clin Pharmacol Ther* **81**, 521-8 (2007).
- (26) Ormeci, A. *et al.* Can *Helicobacter pylori* be eradicated with high-dose proton pump inhibitor in extensive metabolizers with the CYP2C19 genotypic polymorphism? *Eur Rev Med Pharmacol Sci* **20**, 1795-7 (2016).
- (27) Grabowski, B. & Lee, R.D. Absorption, distribution, metabolism and excretion of [14C]dexlansoprazole in healthy male subjects. *Clin Drug Investig* **32**, 319-32 (2012).
- (28) Shakhnovich, V., Ward, R.M. & Kearns, G.L. Failure of proton pump inhibitors to treat GERD in neonates and infants: a question of drug, diagnosis, or design. *Clin Pharmacol Ther* **92**, 388-92 (2012).
- (29) Knebel, W., Tammara, B., Udata, C., Comer, G., Gastonguay, M.R. & Meng, X. Population pharmacokinetic modeling of pantoprazole in pediatric patients from birth to 16 years. *J Clin Pharmacol* **51**, 333-45 (2011).
- (30) Shakhnovich, V. *et al.* A Population-Based Pharmacokinetic Model Approach to Pantoprazole Dosing for Obese Children and Adolescents. *Paediatr Drugs* **20**, 483-95 (2018).
- (31) Kearns, G.L. *et al.* Single-dose pharmacokinetics of oral and intravenous pantoprazole in children and adolescents. *J Clin Pharmacol* **48**, 1356-65 (2008).
- (32) Franciosi, J.P. *et al.* Association Between CYP2C19*17 Alleles and pH Probe Testing Outcomes in Children With Symptomatic Gastroesophageal Reflux. *J Clin Pharmacol* **58**, 89-96 (2018).
- (33) Franciosi, J.P. *et al.* Association between CYP2C19 extensive metabolizer phenotype and childhood anti-reflux surgery following failed proton pump inhibitor medication treatment. *Eur J Pediatr* **177**, 69-77 (2018).
- (34) Mougey, E.B. *et al.* CYP2C19 and STAT6 Variants Influence the Outcome of Proton Pump Inhibitor Therapy in Pediatric Eosinophilic Esophagitis. *J Pediatr Gastroenterol Nutr* **69**, 581-7 (2019).
- (35) Lima, J.J. *et al.* Association of CYP2C19 polymorphisms and lansoprazole-associated respiratory adverse effects in children. *J Pediatr* **163**, 686-91 (2013).
- (36) Cicali, E.J. *et al.* Novel Implementation of Genotype-Guided Proton Pump Inhibitor Medication Therapy in Children: A Pilot, Randomized, Multisite Pragmatic Trial. *Clin Transl Sci* **12**, 172-9 (2019).
- (37) Tang, M. *et al.* Genotype tailored treatment of mild symptomatic acid reflux in children with uncontrolled asthma (GenARA): Rationale and methods. *Contemp Clin Trials* **78**, 27-33 (2019).
- (38) Desta, Z., Zhao, X., Shin, J.G. & Flockhart, D.A. Clinical significance of the cytochrome P450 2C19 genetic polymorphism. *Clin Pharmacokinet* **41**, 913-58 (2002).
- (39) Botton, M.R. *et al.* Structural variation at the CYP2C locus: Characterization of deletion and duplication alleles. *Hum Mutat* **40**, e37-e51 (2019).

Supplement File Title: Supplemental Material

TABLE 1. ASSIGNMENT OF PREDICTED CYP2C19 PHENOTYPE BASED ON GENOTYPE

Predicted phenotype	Genotype	Examples of <i>CYP2C19</i> diplotypes^a
CYP2C19 ultrarapid metabolizer	An individual carrying two increased function alleles	*17/*17
CYP2C19 rapid metabolizer	An individual carrying one normal function allele and one increased function allele	*1/*17
CYP2C19 normal metabolizer	An individual carrying two normal function alleles	*1/*1
CYP2C19 likely intermediate metabolizer ^b	An individual carrying one normal function allele and one decreased function allele or one increased function allele and one decreased function allele or two decreased function alleles	*1/*9, *9/*17, *9/*9
CYP2C19 intermediate metabolizer	An individual carrying one normal function allele and one no function allele or one increased function allele and one no function allele	*1/*2, *1/*3, *2/*17, *3/*17
CYP2C19 likely poor metabolizer ^b	An individual carrying one decreased function allele and one no function allele	*2/*9, *3/*9
CYP2C19 poor metabolizer	An individual carrying two no function alleles	*2/*2, *3/*3, *2/*3
Indeterminate	An individual carrying one or two uncertain function alleles	*1/*12, *2/*12, *12/*14

^aPlease refer to the ***CYP2C19* Diplotype-Phenotype Table** online for a complete list (3, 4).

^bThere are limited data to characterize the function of decreased function alleles.

TABLE 2. DOSING RECOMMENDATIONS FOR OMEPRAZOLE, LANSOPRAZOLE, PANTOPRAZOLE, AND DEXLANSOPRAZOLE BASED ON CYP2C19 PHENOTYPE

CYP2C19 phenotype^a	Implications for phenotypic measures	Therapeutic recommendation	Classification of recommendation^b – omeprazole, lansoprazole, and pantoprazole	Classification of recommendation^b – dexlansoprazole
CYP2C19 ultrarapid metabolizer	Decreased plasma concentrations of PPIs compared to CYP2C19 NMs; increased risk of therapeutic failure	Increase starting daily dose by 100%. Daily dose may be given in divided doses. Monitor for efficacy.	Optional	Optional
CYP2C19 rapid metabolizer	Decreased plasma concentrations of PPIs compared to CYP2C19 NMs; increased risk of therapeutic failure	Initiate standard starting daily dose. Consider increasing dose by 50-100% for the treatment of <i>H. pylori</i> infection and erosive esophagitis. Daily dose may be given in divided doses. Monitor for efficacy.	Moderate	Optional
CYP2C19 normal metabolizer	Normal PPI metabolism; may be at increased risk of therapeutic failure compared to CYP2C19 IMs and PMs	Initiate standard starting daily dose. Consider increasing dose by 50-100% for the treatment of <i>H. pylori</i> infection and erosive esophagitis. Daily dose may be given in divided doses. Monitor for efficacy.	Moderate	Optional
CYP2C19 likely	Likely increased	Initiate standard starting daily dose.	Optional ^c	Optional ^c

intermediate metabolizer	plasma concentration of PPI compared to CYP2C19 NMs; likely increased chance of efficacy and potentially toxicity	For chronic therapy (>12 weeks) and efficacy achieved, consider 50% reduction in daily dose and monitor for continued efficacy.		
CYP2C19 intermediate metabolizer	Increased plasma concentration of PPI compared to CYP2C19 NMs; increased chance of efficacy and potentially toxicity	Initiate standard starting daily dose. For chronic therapy (>12 weeks) and efficacy achieved, consider 50% reduction in daily dose and monitor for continued efficacy.	Optional	Optional
CYP2C19 likely poor metabolizer	Likely increased plasma concentration of PPI compared to CYP2C19 NMs; likely increased chance of efficacy and potentially toxicity	Initiate standard starting daily dose. For chronic therapy (>12 weeks) and efficacy achieved, consider 50% reduction in daily dose and monitor for continued efficacy.	Moderate ^c	Optional ^c
CYP2C19 poor metabolizer	Increased plasma concentration of PPI compared to CYP2C19 NMs; increased chance of efficacy and	Initiate standard starting daily dose. For chronic therapy (>12 weeks) and efficacy achieved, consider 50% reduction in daily dose and monitor	Moderate	Optional

	potentially toxicity	for continued efficacy.		
--	-------------------------	----------------------------	--	--

IM, intermediate metabolizer; NM, normal metabolizer; PM, poor metabolizer; PPI, proton pump inhibitor

^aThe online ***CYP2C19* Frequency Table** provides phenotype frequencies for major race/ethnic groups, and the online ***CYP2C19* Diplotype-Phenotype Table** provides a complete list of possible diplotypes and phenotype assignments (3, 4).

^bRating scheme described in the **Supplemental Material** online.

^cThe strength of recommendation for “likely” phenotypes are the same as their respective confirmed phenotypes. “Likely” indicates the uncertainty in the phenotype assignment, but it is reasonable to apply the recommendation for the confirmed phenotype to the corresponding “likely” phenotype.